

AD-A206 033

DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1b. RESTRICTIVE MARKING

DTIC FILE (X)

2a. SECURITY CLASSIFICATION AUTHORITY

2b. DECLASSIFICATION / DOWNGRADING SCHEDULE

3. DISTRIBUTION / AVAILABILITY OF ABSTRACT
This document has been approved
for public release and no limit
distribution is unlimited.

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

USARIEM, Natick, MA

5. MONITORING ORGANIZATION REPORT NUMBER(S)

6a. NAME OF PERFORMING ORGANIZATION

USARIEM, Natick, MA

6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

6c. ADDRESS (City, State, and ZIP Code)

Natick, MA 01760-5007

7b. ADDRESS (City, State, and ZIP Code)

8a. NAME OF FUNDING / SPONSORING
ORGANIZATION8b. OFFICE SYMBOL
(If applicable)

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

8c. ADDRESS (City, State, and ZIP Code)

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NO.PROJECT
NO.TASK
NO.WORK UNIT
ACCESSION NO.

11. TITLE (Include Security Classification)

(U) Application of a model of exertional heatstroke pathophysiology to cocaine
intoxication.

12. PERSONAL AUTHOR(S)

LE Armstrong, RW Hubbard

13a. TYPE OF REPORT

manuscript

13b. TIME COVERED

FROM Jan 89 TO Jan 89

14. DATE OF REPORT (Year, Month, Day)

11 Jan 1989

15. PAGE COUNT

4

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD	GROUP	SUB-GROUP

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

heatstroke pathophysiology, cocaine intoxication, heat
illness modelling, clinical indices of illness

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

This letter to the editor involves a novel application of The Energy Depletion Model of heatstroke pathophysiology, developed two years ago in the Heat Research Division, USARIEM. This application is made to cocaine intoxication, as reported by Roth et al. (NEJM 319:673-677, 1988) in a recent publication. The Energy Depletion Model is briefly described, five salient reasons are given in support of this application to cocaine intoxication, and future research implications are presented.

DTIC
ELECTE
31 MAR 1989
S E D

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Lawrence E. Armstrong, Ph.D.

22b. TELEPHONE (Include Area Code)

508-651-4873

22c. OFFICE SYMBOL

SGRD-UE-HR

89 3 30 050

Application of a model of exertional heatstroke pathophysiology to cocaine intoxication

To the Editor, New England Journal of Medicine:

We would like to propose a novel cellular model of cocaine intoxication which originates from work on heatstroke pathophysiology^{1,2}. This model emphasizes the role of heat, hypohydration (hypernatremia), and increased neuromuscular activity (i.e. accelerated energy consumption, ion flux, and heat production) in activating the sodium pump. A vicious circle is described which leads to increased intracellular sodium concentration, cell swelling, hyperthermia, energy depletion, metabolic cascades, and irreversible cell damage. Named The Energy Depletion Model, it shifts emphasis from hyperthermia alone to a potential cellular mechanism implicating anaerobic glycolysis and lacticidosis as consequences of the total energy drain on the cell.

For the following reasons, we propose that this model is applicable to cocaine intoxication. (1) Exertional heatstroke^{2,3,9} and cocaine intoxication^{4,5} share many of the same symptoms (Table 1), strongly implicating common pathways during the evolution of each disorder. The symptoms of both illnesses (e.g. elevated serum ALT, AST, CPK; myoglobinuria) suggest increased ion flux, membrane permeability to macromolecules, and thereby a potential increase in permeability to smaller molecular species such as monovalent and divalent cations. (2) We believe that many of the systemic consequences associated with heatstroke have their origin in conditions leading to increased cell permeability and energy depletion². If the commonality of the symptoms between heatstroke and cocaine intoxication involves similar events at this level of structure and function, then the many applications of heat illness modelling could provide new avenues of research for studying cocaine abuse in a performance-based animal model¹. (3) By definition, muscular exercise is an important factor in exertional heatstroke^{1,2,3,9}. Similarly, myoclonus, status

epilepticus, and excessive heat production have been reported following pharmacologic doses of cocaine, amphetamine, phencyclidine (PCP), alcohol, and lithium^{4,5,7}. These drug-induced states result in many of the symptoms^{5,6,7,8} described in Table 1, and should be included among the differential diagnoses^{3,5} in cases of suspected cocaine intoxication and exertional heatstroke. (4) The non-pyrogenic hyperthermia ($> 40^{\circ}\text{C}$) characteristic of cocaine intoxication⁴ (Table 1) agrees well with this model. (5) This model predicts a decreased serum calcium level⁴ (Table 1), theoretically the result of membrane depolarization, dissipation of the sodium gradient, and ATP depletion, which facilitate calcium sequestration within the cell.

Although we have no plans to directly investigate or model cocaine intoxication, we welcome correspondence regarding The Energy Depletion Model^{1,2} from those who do.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<i>form 50 per</i>
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Lawrence E. Armstrong, Ph.D.

Research Physiologist

Roger W. Hubbard, Ph.D.

Director, Heat Research Division

U.S. Army Research Institute of

Environmental Medicine

Kansas Street

Natick, MA 01760-5007

(508) 651-4873



Table 1 - Selected signs and symptoms which may be observed in severe cases of heatstroke^{2,3,9} and cocaine intoxication^{4,5}.

<u>Signs and Symptoms</u>	<u>Cocaine intoxication</u>	<u>Heatstroke</u>
combative behavior	+	+
coma	+	+
nonpyrogenic hyperthermia (> 40°C)	+	+
acute renal failure	+	+
oliguria (< 400ml/24hr)	+	+
hyperuricemia	+	+
rhabdomyolysis	+	+
hypotension	+	+
seizure	+	+
myocardial and/or hepatic damage	+	+
elevated serum ALT, AST, CPK, bilirubin	+	+
myoglobinuria	+	+
serum hypocalcemia	+	+
disseminated intravascular coagulation	+	+
multi-organ hemorrhage	+	+

References

1. Hubbard RW, Matthew CB, Durkot MJ, Francesconi RP. Novel approach to the pathophysiology of heatstroke: The Energy Depletion Model. *Ann Emerg Med* 1987; 16:1066-1075.
2. Hubbard RW, Armstrong LE. The heat illnesses: biochemical, ultrastructural and fluid-electrolyte considerations. In: Pandolf KB, Sawka MN, Gonzalez RR, ed. *Human Performance Physiology and Environmental Medicine at Terrestrial Extremes*. Indianapolis, IN: Benchmark Press, 1988:305-359.
3. Knochel JP. Environmental heat illness. *Arch Intern Med* 1974; 138:841-864.
4. Roth D, Alarcon FJ, Fernandez JA, Preston RA, Bourgoignie JJ. Acute rhabdomyolysis associated with cocaine intoxication. *N Engl J Med* 1988; 319:673-677.
5. Olson KR, Benowitz NL. Environmental and drug-induced hyperthermia. *Emerg Med Clin N Am* 1984; 2:459-474.
6. Granoff AL, Davis JM. Heat illness syndrome and lithium intoxication. *J Clin Psychiatry* 1978; 39:103-107.
7. Kendrick WC, Hull AR, Knochel JP. Rhabdomyolysis and shock after intravenous amphetamine administration. *Ann Intern Med* 1977; 86:381-387.
8. Isner JM, Estes NAM, Thompson PD, Constanzo-Nordin MR, Subramanian R, Miller G, Katsas G, Sweeney K, Sturner WQ. Acute cardiac events temporally related to cocaine abuse. *N Engl J Med* 1986; 315:1438-1443.
9. Shibolet S, Lancaster MC, Danon Y. Heatstroke: a review. *Aviat Space Environ Med* 1976; 47:280-301.